

Geology

Restoration of Laramide right-lateral strike slip in northern New Mexico by using Proterozoic piercing points: Tectonic implications from the Proterozoic to the Cenozoic

K. E. Karlstrom and C. G. Daniel

Geology 1993;21;1139-1142

doi: 10.1130/0091-7613(1993)021<1139:ROLRLS>2.3.CO;2

Email alerting services

click www.gsapubs.org/cgi/alerts to receive free e-mail alerts when new articles cite this article

Subscribe

click www.gsapubs.org/subscriptions/ to subscribe to *Geology*

Permission request

click <http://www.geosociety.org/pubs/copyrt.htm#gsa> to contact GSA

Copyright not claimed on content prepared wholly by U.S. government employees within scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in subsequent works and to make unlimited copies of items in GSA's journals for noncommercial use in classrooms to further education and science. This file may not be posted to any Web site, but authors may post the abstracts only of their articles on their own or their organization's Web site providing the posting includes a reference to the article's full citation. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

Notes

Restoration of Laramide right-lateral strike slip in northern New Mexico by using Proterozoic piercing points: Tectonic implications from the Proterozoic to the Cenozoic

K. E. Karlstrom } Department of Earth and Planetary Sciences, University of New Mexico,
C. G. Daniel } Albuquerque, New Mexico 87131

ABSTRACT

Early Proterozoic rocks in northern New Mexico can be used to document 100–170 km of right-lateral slip along a network of north-striking Laramide (Late Cretaceous–early Tertiary) faults that formed the precursor of the Rio Grande rift. Piercing lines are defined by the intersection of a subhorizontal 4 kbar isobaric surface with steeply dipping stratigraphic markers and regional structures. Restoration of slip provides new insight into the nature of Proterozoic orogenic and crustal province boundaries. The Yavapai crustal province (1.75–1.72 Ga juvenile volcanogenic rocks) is imbricated with overlying Hondo Group sedimentary cover in an originally east-trending ductile thrust belt that formed near the southern margin of the Yavapai province. A series of offset magnetic anomalies (Jemez lineament) is interpreted to mark the thrust-buried boundary between older (Yavapai province) and younger (Mazatzal province) crust. Our microplate tectonic model suggests that Laramide strike-slip movement (New Mexico) was kinematically linked with coeval extensional collapse (Arizona) and foreland shortening (Wyoming) during decoupling and northward movement of the rigid Colorado Plateau relative to the undeformed midcontinent. Crustal-scale strike-slip faults in New Mexico were reactivated during later Tertiary extension and continue to localize high heat flow in the Rio Grande rift and northward into Colorado.

INTRODUCTION

Data from Early Proterozoic rocks support and further constrain Chapin and Cather's (1981) model for a Laramide right-slip fault system in New Mexico. Chapin (1983) suggested ~100 km of slip on the basis of the realignment of northeast-trending magnetic anomalies. Our proposed restoration of 100–170 km of slip provides a testable model for the location and magnitude of individual faults of this system. Restoration of Phanerozoic deformation provides insights into the regional structure of Proterozoic basement rocks of the southwest United States and helps constrain a model involving coeval extension, strike slip, and shortening associated with northward movement of a rigid Colorado Plateau block during Late Cretaceous–early Tertiary time.

RESTORATION OF PROTEROZOIC ROCKS IN NEW MEXICO

Early Proterozoic-age rocks are exposed in isolated mountain uplifts in northern New Mexico and southern Colorado (Fig. 1). Both lithostratigraphic (Bauer and Williams, 1989) and terrane (Grambling et al., 1988) subdivisions emphasize the similarities of rock packages in adjacent ranges. This paper extends the hypothesis of Miller et al. (1963), that Proterozoic rocks and structures can be directly correlated from range to range. The strongest evidence for kilometre-scale strike-slip offsets of Proterozoic rocks and structures comes from the Rio Mora, Truchas, Picuris, and Tusas uplifts, which we propose were once part of a continuous Proterozoic thrust belt (inset in Fig. 1). This proposed alignment is based on correlation of stratigraphic markers, alignment of major thrusts and folds, and similarity of metamorphic grade, as discussed below.

Stratigraphic correlations are based on the distinctive Vadito and Hondo groups. These rocks overlie 1.75–1.72 Ga mafic metavolcanic rocks and associated granodiorite plutons that form the heterogeneous arc-related crust of the southwest United States (Karlstrom and Bowring, 1993). The Vadito Group (1.7 Ga) is dominated by felsic metavolcanic rocks, with a distinctive Mn-rich layer (1–50 m thick) at its top (perhaps a paleosol) that is characterized by Mn-andalusite, piemontite, and other Mn-rich minerals (Grambling and Williams, 1985). This layer can be matched from range to range (Fig. 1). The overlying Hondo Group is dominated by the 1-km-thick Al-rich Ortega quartzite that also correlates from range to range. This quartzite is interpreted as a shallow marine “shelf” sequence derived from the north (Soegaard and Erikson, 1986). The Uncompahgre Formation of the Needle Mountains is similar in thickness and lithology to the Hondo Group and also contains Mn-andalusite (Burns et al., 1980). Major structures also correlate between uplifts. Each range contains two major Proterozoic thrusts (e.g., the Plomo and Pilar faults of the Picuris Mountains; Bauer, 1993) that bound the southernmost of a series of kilometre-scale overturned synclines involving the Hondo Group (Fig. 1; La Madera syncline in the Tusas Mountains; Williams, 1991a; Hondo syncline in the Picuris Mountains, Bauer, 1993).


Direct strike-slip realignment of stratigraphic sequences and structures (Fig. 1) is possible because each range was at about the same crustal depth (12 ± 2 km) after Proterozoic deformation. Metamorphic isograds in New Mexico are subhorizontal and overprint map-scale folds and faults, indicating that metamorphism outlasted or postdated shortening (Grambling et al., 1989). Metamorphic pressure-temperature estimates (Grambling et al., 1989) indicate conditions of 525 ± 25 °C and 3.8–4.5 kbar in each of the ranges. The timing of metamorphism remains controversial (Williams, 1991b), but is bracketed between 1.65 and 1.4 Ga. Thus, the intersection of regionally subhorizontal isobaric surfaces with preexisting subvertical markers and structures creates regional piercing lines that can be used to restore the net (post-1.4 Ga) displacement. This restoration suggests that postmetamorphic vertical displacement and tilting between ranges was insignificant relative to the strike-slip displacement.

A band of northeast-trending magnetic highs (Zietz, 1982) also shows right-lateral offset across north-striking faults (Fig. 1). This band corresponds to the Jemez lineament, an alignment of Cenozoic volcanic centers (Aldrich et al., 1986). The north edge of this zone (Fig. 1) coincides with the southernmost outcrops of Yavapai volcanogenic basement (>1.7 Ga) and Hondo Group cover and the northernmost limit of 1.65 Ga supracrustal rocks. Thus, as in Arizona, the Jemez lineament appears to mark a crustal province boundary.

DISCUSSION OF RIGHT-SLIP FAULTS

Proterozoic exposures in New Mexico are bounded by high-angle faults with complicated movement histories. Our estimates of right-lateral offset since 1.4 Ga include ~80 km of slip across the

Figure 1. Proterozoic rocks of northern New Mexico and southern Colorado. Major range-bounding faults with inferred Late Cretaceous–early Tertiary right slip are shown. Inset shows proposed realignment of Proterozoic rocks in Tusas, Cerro Azul, Picuris, Truchas, and Rio Mora uplifts.



Nacimiento and western rift faults, 15 km across the Tusas-Picuris fault, 37 km across the Picuris-Pecos fault (Miller et al., 1963), and several to tens of kilometres across the Front Range faults (Fig. 1). The best determination for the amount of strike-slip displacement (~50 km) is from the Tusas Mountains east to the Rio Mora uplift (inset in Fig. 1). In addition, Baltz (1967) mapped a series of en echelon folds in Mesozoic and early Tertiary strata west of the Nacimiento fault and estimated ~5 km of Eocene right slip. However, we add 70–80 km of right slip on both this and a concealed western rift fault to accommodate the suggestion by Laughlin (1991) that the Sandia Granite “correlates” with the Fenton Hill granodiorite encountered in drill holes beneath the Jemez Mountains some

80 km north of the Sandia Mountains and to match up the offset magnetic anomalies. Front Range faults show a dextral component, but the magnitude of slip is not well known (Baltz and O'Neill, 1980).

Timing of movement(s) on these faults is not well constrained and probably involved various combinations of late Paleozoic, Laramide, and late Tertiary slip. Miller et al. (1963) also postulated a Precambrian ancestry for the Picuris-Pecos fault on the basis of ductile rotation of foliation into the fault. However, our observations are that both the Picuris-Pecos and Borrego faults are brittle strike-slip features that abruptly truncate Proterozoic foliations. Pennsylvanian movement was dominantly dip-slip (Miller et al., 1963; Soegaard, 1990), as was late Tertiary slip. Thus, following Chapin and

Cather (1981) we propose that most of the right-lateral strike-slip occurred in the Laramide.

IMPLICATIONS FOR PROTEROZOIC CRUSTAL PROVINCES

Controversy regarding the nature and location of both Proterozoic crustal province boundaries and Proterozoic orogenic belts (Karlstrom and Bowring, 1993) is clarified by the realignment of Proterozoic rocks and structures in northern New Mexico (Fig. 2). As in most orogens, deformational provinces do not correspond to crustal provinces, because deformation is transmitted far inboard of the margins of colliding terranes. Our reconstruction (Fig. 3) shows a major zone of tectonic imbrication of Yavapai province basement (>1.7 Ga), Hondo Group cover (1.7 Ga) and Mazatzal province (~ 1.65 Ga) crust in northern New Mexico and southern Colorado. The northern part of this belt (the Mazatzal thrust belt) shows thrust and fold geometries (Williams, 1991a) similar to foreland thrust belts and marks the zone of exposure of thick quartzite successions. It is interpreted to record deformation of the Yavapai province foreland due to the 1.65 Ga collision of the Mazatzal province. A thrust buried crustal province boundary marking the southern limit of known Yavapai crust and the northern limit of 1.65 Ga supracrustal rocks appears to correspond with the northern boundary of the Jemez lineament (Figs. 2 and 3).




Figure 2. Proterozoic rocks of southwest United States and proposed Proterozoic boundaries. Crustal provinces represent blocks of distinct age and composition: Mojave (Archean component), Yavapai (1.75–1.70 Ga juvenile crust), and Mazatzal (1.66–1.60 Ga juvenile crust). Orogens are defined on basis of deformational fronts: Cordilleran orogen (Mesozoic thrusting), Yavapai orogen (1.74–1.69 Ga deformation), Mazatzal orogen (1.65–1.60 Ga deformation), and Grenville orogen (1.3–1.0 Ga deformation).

The east trend of the reconstructed Mazatzal thrust belt in New Mexico is a departure from the northeast trend of tectonic boundaries in Arizona. This bend in the thrust belt may reflect the geometry of the southern continental margin of 1.7 Ga Laurentia (Fig. 3). Such a plate margin configuration may help explain variations in the style and intensity of 1.65 Ga deformation. The 1.7 Ga craton margin in northern New Mexico may have been a salient and thus more intensely deformed, overthickened, and subsequently uplifted by a colliding Mazatzal province. This is compatible with the higher pressure of early Proterozoic-age rocks in New Mexico (4 kbar; Grambling et al., 1989) vs. Arizona (3 kbar; Williams, 1991b) and the wide zone of 1.65 Ga deformation in New Mexico and Colorado.

IMPLICATIONS FOR LARAMIDE MICROPLATE TECTONICS

Our model (Fig. 3) suggests that 100–170 km of Laramide right slip in New Mexico accommodated the northward translation of the relatively rigid Colorado Plateau. Although the details differ, we follow Hamilton (1981) and Chapin and Cather (1981) in trying to link displacements around the Plateau into a kinematically balanced microplate tectonic framework. The linkage between strike slip in New Mexico and Colorado and Laramide foreland shortening in Wyo-




Figure 3. Restoration of Colorado Plateau to its inferred Late Cretaceous position, and kinematic framework of Late Cretaceous–early Tertiary deformation. From 100 to 170 km of northward movement of rigid Colorado Plateau (and Arizona transition zone) produced foreland shortening (Wyoming, Utah, Colorado), strike-slip (New Mexico and western Arizona), and gravitationally induced orogenic collapse (Arizona–Sonora).

ming seems likely because of the continuity of regional fault systems. The timing of movement (Late Cretaceous–early Tertiary) and amount of shortening (~100 km; Gries, 1983) across Wyoming may be comparable to the proposed right slip in New Mexico. The predicted Laramide extension in southern Arizona and left slip (and shortening) across the Hurricane, Grand Wash, and Lemington faults on the western edge of the Colorado Plateau (Fig. 3) are more speculative because of intense late Tertiary deformational overprints. However, there is increasing evidence for Mesozoic–early Tertiary mid-crustal extension in southern California (Foster et al., 1992; Hodges and Walker, 1992) and Arizona (Livaccari, 1991) and for a Laramide ancestry for major structures in the western Colorado Plateau.

The dynamic framework for the northward translation of the plateau is speculative. Our restoration (Fig. 3) shows the Colorado Plateau and Arizona's transition zone structurally above the Tertiary core complexes in the Late Cretaceous. This is compatible with a proposed zone of overthickened crust in southern Arizona (Coney and Harms, 1984; Gastil et al., 1992). Although lower plates of core complexes were finally exhumed to upper crustal levels, and reversal of drainage from northward to southward in central Arizona (Gastil et al., 1992) took place in the middle to late Tertiary, our model follows Livaccari (1991) in suggesting that some (perhaps substantial) fraction of extensional thinning of the orogen also took place in the Laramide. This model is broadly analogous to the gravitational collapse of high topography in the Himalayas and the northward translation of the Tibetan Plateau (Hodges et al., 1992). Late Cenozoic tectonics of the southwest United States were profoundly influenced by this Laramide history in that mid-Tertiary extension in the core complexes was superimposed on crust that had undergone an interplay of contraction and extension in the Late Cretaceous–early Tertiary. Further, the Rio Grande rift (and perhaps the Colorado River corridor) was localized along earlier faults. The Rio Grande rift structures continue to localize high heat flow in a north-trending zone in New Mexico and Colorado (Decker et al., 1988).

ACKNOWLEDGMENTS

We thank numerous people for discussions that led to the refinement of this paper: Paul Bauer, Sue Beard, Steve Cather, Bill Dickinson, Patricia Dickerson, Ernie Duebendorfer, Carol Frost, John Geissman, Gordon Haxel, Rick Livaccari, Bill Muelberger and Mike Williams.

REFERENCES CITED

- Aldrich, M.J., Jr., Laughlin, A.W., Meade, J.S., and Pierce, H.W., 1986, The Jemez lineament: Structural boundaries and control on sedimentary facies, tectonism, and mineralization, in Aldrich, M.J., and Laughlin, A.W., eds., *Proceedings, International Conference on Basement Tectonics*, 6th: Salt Lake City, Utah, International Basement Tectonics Association, p. 104–113.
- Baltz, E.H., 1967, Stratigraphy and regional tectonic implications of part of Upper Cretaceous and Tertiary rocks, east-central San Juan basin, New Mexico: U.S. Geological Survey Professional Paper 552, 101 p.
- Baltz, E.H., and O'Neill, J.M., 1980, Preliminary geologic map of the Mora River area, Sangre de Cristo Mountains, Mora County, New Mexico: U.S. Geological Survey Open-File Report 80-374, scale 1:24,000.
- Bauer, P.W., 1993, Stratigraphic and structural evolution of Proterozoic rocks in the Picuris Range, northern New Mexico: *Journal of Geology*, v. 101, p. 483–500.
- Bauer, P.W., and Williams, M.L., 1989, Stratigraphic nomenclature of Proterozoic rocks, northern New Mexico: Revisions, redefinitions and formalization: *New Mexico Geology*, v. 11, p. 45–52.
- Burns, L.K., Ethridge, F.G., and Tyler, N., 1980, Geology and uranium evaluation of the Precambrian quartz-pebble conglomerates of the Needle Mountains, southwestern Colorado: U.S. Department of Energy Report GJBK-118-80, 161 p.
- Chapin, C.E., 1983, An overview of Laramide wrench faulting in the southern Rocky Mountains with emphasis on petroleum exploration, in Lowell, J.D., ed., *Rocky Mountain forelands and uplifts*: Denver, Colorado, Rocky Mountain Association of Geologists, p. 169–180.
- Chapin, C.E., and Cather, S.M., 1981, Eocene tectonics and sedimentation in the Colorado Plateau–Rocky Mountain area, in Dickenson, W.R., and Payne, M.D., eds., *Relations of tectonics to ore deposits in the southern Cordillera*: Arizona Geological Society Digest, v. 14, p. 33–55.
- Coney, P.J., and Harms, T.A., 1984, Cordilleran metamorphic core complexes: Cenozoic extensional relics of Mesozoic compression: *Geology*, v. 12, p. 550–554.
- Decker, E.R., Heasler, H.P., Buelow, K.L., Baker, K.H., and Hallin, J.S., 1988, Significance of past and recent heat-flow and radioactivity studies in the Southern Rocky Mountains region: *Geological Society of America Bulletin*, v. 100, p. 1851–1885.
- Foster, D.A., Miller, C.A., Harrison, T.M., and Hoisch, T.D., 1992, $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronology and thermobarometry of metamorphism, plutonism, and tectonic denudation in the Old Woman Mountains area, California: *Geological Society of America Bulletin*, v. 104, p. 176–191.
- Gastil, G., Wracher, M., Strand, G., Kear, L.L., Eley, D., Chapman, D., and Anderson, C., 1992, The tectonic history of the southwestern United States and Sonora, Mexico, during the past 100 m.y.: *Tectonics*, v. 11, p. 990–997.
- Grambling, J.A., and Williams, M.L., 1985, The effects of Fe^{3+} and Mn^{3+} on aluminum silicate phase relationships in north-central New Mexico, U.S.A.: *Journal of Petrology*, v. 26, p. 324–354.
- Grambling, J.A., Williams, M.L., and Mawer, C.K., 1988, The Proterozoic tectonic assembly of New Mexico: *Geology*, v. 16, p. 724–727.
- Grambling, J.A., Williams, M.L., Smith, R.F., and Mawer, C.K., 1989, The role of crustal extension in the metamorphism of Proterozoic rocks in New Mexico, in Grambling, J.A., and Tewksbury, B.J., eds., *Proterozoic geology of the Southern Rocky Mountains*: Geological Society of America Special Paper 235, p. 87–110.
- Gries, R., 1983, North-south compression of Rocky Mountain foreland structures, in Lowell, J.D., ed., *Rocky Mountain forelands and uplifts*: Denver, Colorado, Rocky Mountain Association of Geologists, p. 9–32.
- Hamilton, W., 1981, Plate-tectonic mechanism of Laramide deformation, in Boyd, D.W., and Lillegraven, J.A., eds., *Rocky Mountain foreland basement tectonics: Contributions to Geology*, v. 19, p. 87–92.
- Hodges, K.V., and Walker, J.D., 1992, Extension in the Cretaceous Sevier orogen, North American Cordillera: *Geological Society of America Bulletin*, v. 104, p. 560–569.
- Hodges, K.V., Parrish, R., Housh, T., Lux, D., Burchfiel, B.C., Royden, L., and Chen, Z., 1992, Simultaneous Miocene extension and shortening in the Himalayan Orogen: *Science*, v. 258, p. 1466–1470.
- Karlstrom, K.E., and Bowring, S.A., 1993, Proterozoic orogenic history of Arizona, in Van Schmus, R.A., and Bickford, M.E., eds., *Precambrian continental United States*: Boulder, Colorado, Geological Society of America, *Geology of North America*, v. C-2, p. 188–211.
- Laughlin, A.W., 1991, Fenton Hill granodiorite—An 80 km (50 mi) right-lateral offset of the Sandia pluton?: *New Mexico Geology*, v. 13, p. 55–59.
- Livaccari, R.F., 1991, Role of crustal thickening and extensional collapse in the tectonic evolution of the Sevier-Laramide orogeny, western United States: *Geology*, v. 19, p. 1104–1107.
- Miller, J.P., Montgomery, A., and Sutherland, P.K., 1963, *Geology of part of the Sangre de Cristo Mountains, New Mexico*: New Mexico Bureau of Mines and Mineral Resources Memoir 11, 106 p.
- Soegaard, K., 1990, Fan-delta and braid-delta systems in Pennsylvanian Sandia Formation, Taos Trough, northern New Mexico: Depositional and tectonic features: *Geological Society of America Bulletin*, v. 102, p. 1325–1343.
- Soegaard, K., and Eriksson, K.A., 1986, Transition from arc volcanism to stable-shelf and subsequent convergent-margin sedimentation in northern New Mexico from 1.76 Ga: *Journal of Geology*, v. 94, p. 47–66.
- Williams, M.L., 1991a, Heterogeneous deformation in a ductile fold-thrust belt: The Proterozoic structural history of the Tusas Mountains, New Mexico: *Geological Society of America Bulletin*, v. 103, p. 171–188.
- Williams, M.L., 1991b, Overview of Proterozoic metamorphism in Arizona: *Arizona Geological Society Digest*, v. 19, p. 11–26.
- Zietz, Isodore, 1982, Composite magnetic anomaly map of the United States: U.S. Geological Survey Geophysical Investigations Map GP-954-A, scale 1:2,500,000.

Manuscript received April 30, 1993

Revised manuscript received September 7, 1993

Manuscript accepted September 13, 1993